

Lodi Memorial Hospital  
975 S. Fairmont Ave.  
Lodi, CA 95240  
(209)334-3411

Patient Name: JOHNSON, HAZEL SHIRLEY  
Age/Sex: 77 F  
Attend MD: Sorour, Nagui N MD

Rm/Bed: 3S-364-  
Adm Date: 05/27/10 at 0943

MR#: M200166  
Acct#: V021432877

ENTERED

Patient Belongings

PATIENT BELONGINGS-----To be completed upon admission, room change and discharge

Received patient from: Emergency Room Does patient have belongings? Y

Glasses? N Disposition-

Contact lenses? N Disposition-

Purse/wallet? N Disposition-

Medications? N Disposition-

Hearing aid(s)? N Disposition-

Dentures? Y Disposition- With patient

Mobility aid(s)? N Disposition-

Prosthesis? N Disposition-

Clothing? Y Disposition-

Side:

Type: Partial upper

Type-

Type-

Type- Sweatpants

Sweatshirt

Jacket

Underwear, socks

Underwear

Slippers

Robe

Type- Necklace

Jewelry? Y Disposition- With patient

Does this inventory match the previous list?

If not please describe action taken

Comment/

Discharged to-

06/02/10 SINGH, BALBIR CNA

Electronically completed by the user listed above.

As a patient, I am encouraged to leave personal items at home. It is my responsibility to inform nursing staff when adding or deleting items from the patient belonging inventory list. The hospital is not liable for the loss or damage to any money, jewelry, documents, or other articles that are not placed in the safe. Hospital liability for loss of any personal property deposited with hospital for safekeeping is limited by law to five hundred (\$500) unless I receive a written receipt for a greater amount from the hospital.

Family representative taking belongings home \_\_\_\_\_ Date \_\_\_\_\_

Patient/Significant Other Signature: \_\_\_\_\_

Staff Signature: \_\_\_\_\_

Transferred to: \_\_\_\_\_ Reconciliation Validated by \_\_\_\_\_ Date \_\_\_\_\_

# Lexi-Patient Education

## Amoxicillin and Clavulanate Potassium

Printed on Wed, June 2, 2010

**U.S. Brand Names** Amoclan; Augmentin ES-600®; Augmentin XR®; Augmentin®

**Canadian Brand Names** Amoxi-Clav; Apo-Amoxi-Clav®; Augmentin®; Clavulin®; Novo-Clavamoxin; ratio-Aclavulanate

**Mexican Brand Names** Acarbixin; Amobay Cl; Amoxiclav; Amoxiclav-BID; Augmentin; Clamoxin; Clavant; Clavipen; Clavucyd; Clavulin; Clavuser; Gimaclav; Moxlin; Riclasip; Servamox; Sinufin

**Pharmacologic Category** Antibiotic, Penicillin

### Reasons not to take this medicine

- If you have an allergy to amoxicillin, clavulanate potassium, any penicillin, or any other part of this medicine.
- Tell healthcare provider if you are allergic to any medicine. Make sure to tell about the allergy and how it affected you. This includes telling about rash; hives; itching; shortness of breath; wheezing; cough; swelling of face, lips, tongue, or throat; or any other symptoms involved.
- If this medicine caused liver problems before.
- Augmentin XR™: If you have severe kidney disease or are on dialysis.

### What is this medicine used for?

- This medicine is used to treat a variety of bacterial infections.

### How does it work?

- Amoxicillin works to injure the bacteria and fight the infection.
- Clavulanate protects amoxicillin from being destroyed by certain bacteria.

### How is it best taken?

- To gain the most benefit, do not miss doses.
- Use prescription as directed, even if feeling better.
- Take this medicine with or without food. Take with food if it causes an upset stomach.
- Chewable tablet: Chew or crush well. Mix crushed tablet with food. Do not swallow whole.
- A liquid (suspension) is available if you cannot swallow pills. Shake well before use.
- Those who have feeding tubes can also use the liquid. Flush the feeding tube before and after medicine is given.
- Scored long-acting product: Do not chew or crush.

- Unscored long-acting product: Swallow whole. Do not chew, break, or crush.

**What do I do if I miss a dose? (does not apply to patients in the hospital)**

- Take a missed dose as soon as possible.
- If it is almost time for the next dose, skip the missed dose and return to your regular schedule.
- Do not take a double dose or extra doses.
- Do not change dose or stop medicine. Talk with healthcare provider.

**What are the precautions when taking this medicine?**

- If you have kidney disease, talk with healthcare provider.
- If you have liver disease, talk with healthcare provider.
- If you have PKU, talk with healthcare provider. Some products do contain phenylalanine.
- Check medicines with healthcare provider. This medicine may not mix well with other medicines.
- Avoid alcohol (includes wine, beer, and liquor).
- Tell healthcare provider if you are pregnant or plan on getting pregnant.
- Tell healthcare provider if you are breast-feeding.

**What are some possible side effects of this medicine?**

- Nausea or vomiting. Small frequent meals, frequent mouth care, sucking hard, sugar-free candy, or chewing sugar-free gum may help.
- Diarrhea. Yogurt, *Bifidobacterium bifidum*, or *Lactobacillus acidophilus* may help. These products are available at health food stores or in some pharmacies.
- For females, vaginal yeast infection. Report itching or discharge.

**What should I monitor?**

- Change in condition being treated. Is it better, worse, or about the same?
- If you are diabetic, this medicine may interfere with Clinitest®. Use another method of urine glucose testing like Clinistix® or Tes-Tape®.

**Reasons to call healthcare provider immediately**

- If you suspect an overdose, call your local poison control center or emergency department immediately.
- Signs of a life-threatening reaction. These include wheezing; chest tightness; fever; itching; bad cough; blue skin color; fits; or swelling of face, lips, tongue, or throat.
- Severe nausea or vomiting.
- Severe diarrhea, even after medicine is stopped.
- Unusual bruising or bleeding.

- Dark urine or yellow skin or eyes.
- Not able to eat.
- Feeling extremely tired or weak.
- Any rash.
- No improvement in condition or feeling worse.

**How should I store this medicine?**

- Store tablets at room temperature.
- Protect tablets from moisture. Do not store in a bathroom or kitchen.
- Store liquid (suspension) in a refrigerator. Do not freeze. Throw away any unused portion after 10 days.

**General statements**

- If you have a life-threatening allergy, wear allergy identification at all times.
- Do not share your medicine with others and do not take anyone else's medicine.
- Keep all medicine out of the reach of children and pets.
- Most medicines can be thrown away in household trash after mixing with coffee grounds or kitty litter and sealing in a plastic bag.
- Keep a list of all your medicines (prescription, natural products, supplements, vitamins, over-the-counter) with you. Give this list to healthcare provider (doctor, nurse, nurse practitioner, pharmacist, physician assistant).
- Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088 or in Canada to Health Canada's Canada Vigilance Program at 1-866-234-2345.
- Talk with healthcare provider before starting any new medicine, including over-the-counter, natural products, or vitamins.

Disclaimer: We want you to get the most benefit you can from your medicine. This patient leaflet is a summary of useful information to help you understand and take it safely. Other information about this medicine may be important for you to know. Please talk with your healthcare provider for more information and your special health needs.