

Advanced care for the heart... from the heart 1801 E. March Lane, Suite D400, Stockton, CA 95210

Should you need to reschedule, please call (209) 46 PHARM ACOLOGICAL MYOVIEW TEST This study is done to evaluate the blood supply to the heart. An intravenous line is required in order to make in injection. One of the medicines Regadence sine, Dipyridamole or Adenosine is infused in order to simulate exercise for the heart's arteries. No physical exercise is required; however, sometimes you may be asked to walk slowly, if possible, on a flat surface. Following this, a small amount of radioactive imaging agent, Myoview, is injected so that im ages, of the heart can be obtained. Images are obtained with a nuclear camera after injection and then repeated at a different time. The test will be performed in a one or two day protocol. The test will take approximately two hours. DO NOT EAT OR DRINK ANYTHING FOR 12 HOURS PRIOR TO ALL VARIETIES OF Int. STRESS PROCEDURES. DO NOT CONSUME ANY CAFFIENE CONTAINING BEVERAGES SUCH AS COFFEE, TEA. COLA, OR CHOCOLATE EVEN IF IT STATES DECAFFIENATED FOR 24 HOURS BEFORE THE TEST. DO NOT TAKE ANACIN OR EXCEDRIN AS THESE CONTAIN CAFFIENE, If there is a possibility that you have a cold, fever or if you are pregnant or nursing please notify the office Females: The tracer that is injected during the procedure has the potential to cause birth defects in fetuses. For this reason, any females under age 50 or who are still having menstrual periods and have not had surgery to prevent pregnancies are required to undergo a pregnancy test prior to the procedure. This warning applies to me and I will undergo pregnancy testing two days prior to my test. This warning does not apply to me since I am over age 50 or I am no longer having menstrual periods or I have had a surgery that prevents pregnancies. If you have emphysema (COPD) or asthma, please bring your inhalers. You may use your inhaler, but please let our stress test rechnician know if you used it or had any wheezing that day prior to the test. DO NOT take asthma medications. Int If you are a diabetic, your test will be scheduled for a morning appointment. Please do not eat breakfast or take your morning diabetes medicine until after the test is completed. We do not anticipate you having a low blood sugar reaction; however, bring something with you to treat a low blood sugar reaction just in case you do experience this. Please wear button-up, short sleeved, shirts/blouses and pants or shorts with flat tennis shoes. No dresses or nylons. (Do not wear clothing with metal buttons or snaps). MEDICATIONS YOU <u>SHOULD NOT TAKE 72 HOURS PRIOR</u> TO YOUR TEST:THEODUR, THEO-24 EXTENDED RELEASE, UNIDUR, UNIPHYL, SLO-PHYLLIN, THEOLAIR, DYPHYLLINE, LUFYLLIN, THEOPHYLLINE. DO NOT TAKE VIAGRA, CIALIS or LEVITRA FOR 7 DAYS PRIOR TO THE TEST. YOUR MEDICATION IS NOT ON THE LISTS ABOVE YOU MAY TAKE IT. PLEASE CALL IF YOU ARE NOT SURE. SPECIAL INSTRUCTIONS: Take all medications as usual, except: Discontinue the following medications 24 hours prior to your test: Discontinue the following medications 48 hours prior to your test: Resume taking these medications upon completion of your stress test. Registration/Prep Time: Test Time: Registration/Prep Time: Test Time: Please do not bring young children with you, since they will not be allowed in the testing area due to radioactivity nor can they be left unattended in the oom. No one other than an interpreter may accompany the patient in the nuclear lab due to radioactive exposure. Kindly give 24-hour notice if you are unable to keep your appointment. The radionucleide imaging agent used is mixed the clay before especially for your body weight and cannot be used on anyone but you. A \$200.00 unused radionucleide imaging agent fee may be charged for all appointments not kept or the above instructions not complied with. This fee is not covered by insurance.





1801 E MARCH LANE STE. D400
 STOCKTON, CA 95210
 (209) 464-3615

(2	(19) 33 4-0651	(209) 464-3615
squeeze a	2D M-MODE ECHOCARDIOGRAM Study where sound waves are used to see the heart and evaluate its function. The size of the procedure requires lying on a cechnician obtains the necessary views of the heart. Please wear a shirt/blouse since you will be	comfortable hed for approximately 30 minutes and
procedure	CAROTID DUPLEX ULTRASOUND study where sound waves are used to see whether there are blockages in the carotid arteries, requires lying on a comfortable bed for approximately 30 minutes while the echo technician ob will be asked to disrobe from the waist up.	which are located on both sides of the neck. This tains the necessary views. Please wear a shirt/blouse
This is a s	VENOUS DUPLEX ULTRASOUND Tudy where sound waves are used to see whether there are blood clots in the deep blood vessels blobed for approximately 30 minutes while the echo technician obtains the necessary views.	located in the legs. This procedure requires lying on
This is a svaricose v	VARICOSE VEIN STUDY study with sound waves that looks for underlying causes of varicose veins in your legs. It caeins in the leg aren't very noticeable. You may be standing during part of the exam. Please expecting in the leg aren't very noticeable.	an sometimes find underlying problems even if the ct the test to take approximately 60 minutes.
This study calculate a	ANKLE / BRACHIAL INDICES (ABI) is performed to see whether there are blockages in the arteries in the legs. It involves taking in index. This will be done at rest and, if necessary, following light exercise on a treadmill. To blood flow in your feet. Please wear flat, rubber soled shoes.	blood pressure readings in your ankles and arms to he technician also may use ultrasound imaging to
This is a st lying on a	ARTERIAL DOPPLER OF THE LOWER EXTREMITIES Tudy where sound waves are used to see whether there are blockages in arteries located in the le comfortable bed for approximately two hours while the echo technician obtains the necessary prior to the test.	egs due to atherosclerosis. This procedure requires views. If you are diabetic, you may eat a small,
This is a str down through the necessar	ABDOMINAL AORTIC ULTRASOUND Lidy where sound waves are used to see whether there are any abnormalities in the wall of the gh your abdomen to your legs. This procedure requires lying on a comfortable bed for approximing views. Please do not chew gum or consume any food that causes bloating or gas for 24 is om midnight the night before the procedure.	ately 30 minutes while the echo technicion alt.
This is a sturequires lying	RENAL DUPLEX ULTRASOUND Ledy where sound waves are used to see whether there are any blockages of the main kidney artering on a comfortable bed for approximately 1 – 2 hours while the echo technician obtains the ray food that causes bloating or gas for 24 hours prior to the exam and do not eat or drink	recessary views Please do not show
This is a stue exercise. Us procedure de plenty of flu	TRESS ECHOCARDIOGRAM addy where a treadmill stress test is performed with the addition of cardiac ultrasound (echocar se of echocardiography with stress testing increases the accuracy of the test. The test will require escribed above, plan to be here approximately 45 minutes. Please wear cool, comfortable, clot aids during the day of the test so you are fully hydrated. Please DO NOT apply lotion to a use an inhaler, please bring it with you.	walking briskly on a treadmill, in addition to the
	NSTRUCTIONS:	
	Take diabetes medications as follows:	
_	Discontinue the following medications 24 hours prior to your test::	
	Resume taking these medications upon completion of your stress	test.
nt.	A \$50.00 missed appointment fee may be charged for appointments not kept. Please numbers above 24 hours in advance if you are unable to keep your scheduled appoint To ensure our technicians are able to provide a high quality study, no children at testing area. Children left unattended in the waiting room must be age 14 or over	call the office at the phone

Patient Name:

Date: 9713

Registration Time:

im

DOB:____

Test Time:

ECHO TESTING ORDER FORM